

Fingolimod Accord

Pregnancy-Specific Patient Reminder Card

Before starting Fingolimod Accord treatment

- **Fingolimod should not be taken if you are pregnant, trying to become pregnant or a woman of child-bearing potential (including female adolescent) not using effective contraception.**
- Fingolimod can harm an unborn baby (the medicine is said to be 'teratogenic'). Before you start treatment with Fingolimod, your doctor will inform you about the risks to an unborn baby and explain what you should do to avoid becoming pregnant while you are taking Fingolimod.
- Your doctor will ask you to take a pregnancy test to ensure that you are not pregnant before you start treatment with Fingolimod.
- Your doctor will explain the need for effective contraception while on treatment and for 2 months after stopping treatment. Talk to your doctor about the most effective contraception options available to you.
- Please read the Fingolimod Patient/Parent/Caregiver Guide provided by your doctor.

While you are taking Fingolimod Accord

- You should not become pregnant while taking Fingolimod and for 2 months after stopping treatment.
- You should use effective contraception while taking Fingolimod and for 2 months after you stop taking it, to avoid becoming pregnant.
- Your doctor will ask you to repeat pregnancy tests at suitable intervals during treatment.
- Your doctor will provide you with regular counselling about Fingolimod's serious risk to an unborn baby.
- Tell your doctor straight away if you become pregnant or if you want to become pregnant because your doctor will stop your treatment with Fingolimod Accord.
- In the event of a pregnancy, Fingolimod should be discontinued. Your doctor will provide counselling on the serious risk to the unborn baby and ultrasonography examinations should be performed.

After stopping Fingolimod Accord treatment

- Inform your doctor immediately if you believe your MS is getting worse (e.g. weakness or visual changes) or if you notice any new symptoms after stopping treatment with Fingolimod due to pregnancy.
- Effective contraception is needed for 2 months after stopping Fingolimod treatment because of the length of time it takes for Fingolimod to leave the body.

Product Registrant:

Accord Healthcare Private Limited
6 Shenton Way, OUE Downtown #38-01
Singapore, 068809

Phone: +65 6714 6700

Website: <https://accord-healthcare.asia/>

This document has been approved by HSA as of 29-Jun-2022.