

I am on anticoagulation treatment with Leaxolar (dabigatran etexilate)

Please complete this section or ask your doctor to do it

Name of the patient

Date of Birth

Indication for Anticoagulation

Dose of Leaxolar

For distribution by healthcare professionals to patients who have been prescribed with Leaxolar

In case of emergency, please contact:

Doctor's Name

Doctor's Contact

Relative's name

Relative's Contact

This document has been approved by HSA on 31 December 2025.

# Leaxolar

dabigatran etexilate

## Patient Alert Card

- Keep this card with you at all times.
- Make sure to use the latest version.
- Your doctor has initiated treatment with Leaxolar. In order to use Leaxolar safely, please consider the important information inside.
- As the patient alert card contains important information about your treatment, please carry this card with you at all times to inform healthcare professionals about your intake of Leaxolar.

## Leaxolar Information for Patients

---

### About your treatment

- Leaxolar thins the blood, which prevents you from getting dangerous blood clots.
- Follow your doctor's instructions when taking Leaxolar. Never skip a dose or stop the intake of Leaxolar without talking to your doctor.
- Inform your doctor about all medicines you are currently taking.
- Inform your doctor about your intake of Leaxolar before any surgery/invasive procedure. Leaxolar has a specific reversal agent (Idarucizumab) that removes the blood thinning effect of Leaxolar in case you need an emergency surgery or have a serious bleed. Please consult your doctor for further information.
- Leaxolar can be taken with or without food. Swallow the capsule whole with a glass of water. Do not break, chew, or empty the contents from the capsule.

### When to seek medical advice

- Taking Leaxolar may increase the risk of bleeding. Speak to your doctor immediately if you experience any of the following possible signs and symptoms of bleeding such as: unusual bruising, nosebleeds, abnormal menstrual flow or vaginal bleeding, blood in the urine, red/black stools.
- If you fall or injure yourself, especially if you hit your head, urgently seek medical advice.
- Do not stop intake of Leaxolar without talking to your doctor, if you experience heartburn, nausea, vomiting, stomach discomfort, bloating or upper abdominal pain.

For further information please contact your doctor or pharmacist

---