doctor's number Date Lormalzi[®] started

Prescribing doctor's name Prescribing

person and number

AUIPU S MANPA

important Contact Information:

I hese are not all the possible side effects of Lormals!". Talk to your doctor right away if you experience any side effects while taking this medication.

(seruzies) eti T 👰

ssəuizzid 🕵



Loss of consciousness















Although most people who developed ARIA do not have symptoms, some people may experience them. **Seek medical** a**stentions** if you experience any of these symptoms.

AIAA to emotqmy2

 Your doctor will arrange brain magnetic resonance imaging Midl8] scarse below your second, find, fourth, and easewith doses of Lormalis*. It is very important that you aftend all your scheduled MRI scan appointments to check if you have ARIA.
 Your doctor may schedule additional scans at other times during treatment if deemed necessary.

 Infrequently, larger areas of bleeding in the brain, known as infracerebral haemorrhage (ICH), can occur.

- Commail/er acuse side ellerica of harm acustiling and bleeding, also known as amyloid-resided imaging abnormalities (ARBA), which can be serious, life-threatening abnormalities (ARBA), which can be serious, life-threatening of the brain, with or withhout small spots of bleeding in or on the surface of the brain.

Amyloid-related Imaging Abnormalities (ARIA) with Lormalzi®

acetytsalicytic acid).

 Inform your doctor about any new medications taken during treatment, including medicines to reduce blood clots professional, especially medicines to reduce blood clots from forming lanti-thrombotic medicines, including

 Keep this card with you or your caregiver, and show it to any healthcare professional involved in your treatment so that they know you are taking Lormals![®].

> Information for Patients and Caregivers:

Important Information for Healthcare Professionals

This patient has received Lormatize Idonanemabl, a monoclonal antibody directed against aggregated forms of beta amyloid, which can cause amyloid-related imaging abnormalities (ARIA). Assess the patient for symptoms of ARIA, including but not limited to those listed on this card. As ARIA can cause focal neurologic deficits that can mimic an ischaemic stroke, please consider it symptoms could be due to ARIA before giving thrombolytic therapy. Please contact the patient's prescribing doctor for further information.

Please report suspected adverse events (AEs) to HSA via their online reporting portal [https://www.hsa.gov.sg/adverse-events] and email a copy of the submitted HSA AE reporting form to hee-py.sin@dksh.com



Please scan the QR code to access the full prescribing information for Lormalzi®.


This document has been approved by HSA as of 11/03/2025

lormalzi (donanemab)

Patient Alert Card

Important Safety Information for patients taking Lormalzi® (donanemab)

Please keep this card with you at all times.



Please keep inis card with you at all times.

Important Safety Information for patients taking Lormatzi®

Patient Alert Card

(qousueusp)

This document has been approved by HSA as of 11/03/2025.

Please scan the QR code to access the full prescribing information for Lormation for



hec-pv.sin@dksh.com.

Please report suspected adverse events (BAS) to ASH vis their online reporting portal (https://www.hsa.go.vop.sg/adverse-events) and it is exporting to ming to ming a ASH baltimus to copy of the submitted that a second or ming to ming to

doctor for further information.

The petient has the ecored Lement's floonnemable, a monocloonal antibody directed against aggregated forms of best amyolid, which can cause amyloid-related imaging abnormalities [ARIA]. Individual of the cause focal limited to hose fusted on this card. As ARIA including but not limited to hose fusted on this card. As ARIA can cause focal immited to hose fusted on this card. As ARIA can cause focal consider if symptoms could be use to ARIA before giving through the considering and the considering and the card of the c

Important Information for Healthcare Professionals

Information for Patients and Caregivers:

- Keep this card with you or your caregiver, and show it to any healthcare professional involved in your treatment so that they know you are taking Lormalzi[®].
- Inform your doctor about any new medications taken during treatment, including medicines supplied by any healthcare professional, especialtly medicines to reduce blood clots from forming lanti-thrombotic medicines, including acet visalicitic acidi).

Amyloid-related Imaging Abnormalities (ARIA) with Lormalzi®

- Lormatzi® can cause side effects of brain swelling and bleeding, also known as amyloid-related imaging abnormalities (ARIA), which can be serious, life-threatening or fatal. This is characterised by temporary swelling in areas of the brain, with or without small spots of bleeding in or on the surface of the brain.
- Infrequently, larger areas of bleeding in the brain, known as intracerebral haemorrhage (ICH), can occur.
- Your doctor will arrange brain magnetic resonance imaging (MRI) scans before your second, third, fourth, and seventh doses of Lormalizi⁸. It is very important that you attend all your scheduled MRI scan appointments to check if you have ARIA. Your doctor may schedule additional scans at other times during treatment if deemed necessary.

Symptoms of ARIA

Although most people who developed ARIA do not have symptoms, some people may experience them. Seek medical attention right away if you experience any of these symptoms.

(Headache

Trer

Confusion

Vision chang

(R) Vomiting

Vision changes

Speech disturbances

Loss of balance

Loss of consciousness

(Dizziness

Fits (seizures)

These are not all the possible side effects of Lormalzi[®]. Talk to your doctor right away if you experience any side effects while taking this medication.

Important Contact Information:

Patient's name :

Emergency contact : person and number

Prescribing : doctor's name

Prescribing doctor's number

Date Lormalzi®