## PATIENT SAFETY INFORMATION CARD

Asccination Date
Tel. number
Pmsin naisisyd
Hospital where treated
emen fneite?
Show this card to any doctor involved in your health care.

Patients receiving SOLIRIS® should carry this card at all times.

.62U2/0U/#I no





AstraZeneca

## Important Safety Information for Patients Taking SOLIRIS® (eculizumab)

SOLIRIS<sup>®</sup> can lower the ability of your immune system to fight infections, especially meningococcal infection, which requires <u>immediate</u> medical attention. If you experience any of the following symptoms, you should <u>immediately call your doctor or seek emergency medical care</u>, preferably in a major emergency medical care center.

- · headache with nausea or vomiting
- · headache with a stiff neck or stiff back
- fever
- rash

- confusion
- muscle aches with flu-like symptoms
- · eyes sensitive to light



Get emergency medical care right away if you have any of these signs or symptoms and  $\underline{show this card}$ .

Even if you stop using SOLIRIS<sup>®</sup>, keep this card with you for 3 months after your last SOLIRIS<sup>®</sup> dose. Your risk of meningococcal infection may continue for several weeks after your last dose of SOLIRIS<sup>®</sup>.

## Information for the Treating Physician



This patient has been prescribed SOLIRIS<sup>®</sup> (eculizumab) therapy, which increases the patient's susceptibility to meningococcal infection (*Neisseria meningitidis*) or other general infections.

- Meningococcal infections may become rapidly life-threatening or fatal if not recognised and treated early.
- Evaluate immediately if infection is suspected and treat with appropriate antibiotics if necessary.
- · Contact prescribing physician (refer to back of card) as soon as possible.

For more information about SOLIRIS®, please refer to the Package Insert (PI) or e-mail: <u>MedInfo.SG@astrazeneca.com</u>.

In case of safety concerns, please call +65 6812 4700 or e-mail: <u>PatientSafety.SG@astrazeneca.com</u>.