

Important Contact Information

Name of TEVIMBRA™ Prescriber:

Phone:

Patient Name:

Patient Phone:

Emergency Contact Name:

Emergency Contact (Phone):

Important Information for Health Care Providers

This patient is being treated with TEVIMBRA™ (tislelizumab), which can cause immune-mediated adverse reactions that may appear at any time during treatment or even after treatment. Assess patients for signs and symptoms of immune-mediated adverse reactions, including but not limited to those listed on this card. Early diagnosis and appropriate management are essential to minimize any consequences of immune-mediated adverse reactions.

For suspected immune-mediated adverse reactions, ensure adequate evaluation to confirm the etiology and exclude other causes. Depending on the type and severity of immune-mediated adverse reactions, consider withholding tislelizumab and administering corticosteroids. Specific guidelines for managing immune-mediated adverse reactions are available in the Singapore Package Insert for tislelizumab. Consultation with an oncologist or other medical specialists may be helpful for management of organ-specific immune-mediated adverse reactions.

All suspected adverse drug reactions (ADRs) can also be reported to HSA via <https://www.hsa.gov.sg/adverse-events> or the Marketing Authorization Holder BeiGene (www.beigene.com/adverse-event-form).



BeiGene

BeiGene Singapore Pte. Ltd.

38 Beach Road, #23-11 South Beach Tower, Singapore 189787

Version 2.0, May 2025

This document has been approved by HSA as of 19-Sep-2024

0225-TEV-PRC-002

PATIENT CARD



TEVIMBRA™ can cause serious side effects, which can sometimes become life-threatening and can lead to death. You may experience side effects in more than one body part at the same time. These side effects may happen anytime during treatment or even after your treatment has ended. Your doctor may provide other medicines to treat your symptoms, delay a next infusion or completely stop further treatment if the side effects are too severe.

Important safety Information

- ▼ If you have any questions about how TEVIMBRA™ works or why this medicine has been prescribed for you, ask your doctor.
- ▼ If you experience any adverse reactions (side effects), talk to your doctor. This includes any possible side effects not listed on this card.
- ▼ Do not attempt to diagnose or self-treat any symptoms without consulting a healthcare professional first.
- ▼ It is important that you carry the **Patient Card** with you at all times and show it to healthcare professionals at all medical visits (e.g., emergency healthcare professionals) to help with diagnosing and treating immune-related adverse reactions.

Look out for serious adverse reactions (side effects)

Tell your doctor **immediately** if you get any of these side effects during treatment with TEVIMBRA™ :

- ▼ **inflammation of the lungs (pneumonitis):** Shortness of breath, chest pain, new or worsening cough.
- ▼ **inflammation of the liver (hepatitis) or other liver problems:** Feeling sick (nausea), vomiting, pain on right side of the stomach area, loss of appetite, yellowing of skin or the whites of eyes, bleeding or bruising more easily than normal, tiredness, dark colored urine.
- ▼ **inflammation of the large bowel (colitis):** Diarrhoea, or more bowel movements than normal, black, tarry, sticky stools, blood or mucus in stools, severe pain or tenderness in the stomach area, feeling sick (nausea), vomiting.
- ▼ **inflammation of the kidney (nephritis):** Changes in the amount and color of the urine, pain while urinating, pain in kidney area, blood in urine, swelling in the ankles, loss of appetite.
- ▼ **severe skin reactions:** Fever, flu-like symptoms, rash, itching, skin blistering or ulcers in the mouth or on other moist surfaces of the nose, throat, genital area.
- ▼ **problems with hormone-producing glands (especially the adrenal, pituitary, or thyroid glands):** Fast heart rate, increased sweating, hyperactivity, extreme tiredness, swelling at the base of the neck, pain in front of the throat, weight gain or weight loss, dizziness or fainting, confusion, hair loss, skin changes, feeling cold, constipation, headaches that will not go away or unusual headaches, decreased blood pressure, vision changes, excessive thirst, changes in mood or behavior, such as decreased sex drive, irritability or forgetfulness, deeper voice.

- ▼ **type 1 diabetes mellitus:** High blood sugar, feeling more hungry or thirsty than normal, passing urine more often than normal, fruity-smelling breath.
- ▼ **infusion-related reaction:** Chills, shaking, itching, rash, flushing, shortness of breath, wheezing, dizziness, fever (may occur during or up to 24 hours after infusion).
- ▼ **inflammation of the muscles (myositis):** Muscle pain, stiffness, weakness, chest pain, severe tiredness.
- ▼ **inflammation of the heart muscle (myocarditis):** Chest pain, rapid or abnormal heartbeat, shortness of breath at rest or during activity, fluid build-up with swelling of the legs, ankles and feet, tiredness.
- ▼ **inflammation of the joints (arthritis):** Joint pain, stiffness, swelling or redness, decreased range of motion in the joints.
- ▼ **inflammatory disorder (polymyalgia rheumatica):** Muscle pain and stiffness, especially in the neck, upper arms, wrists, elbows, and shoulders, knees and hips, buttocks, or thighs.
- ▼ **inflammation of the membrane around the heart (pericarditis):** Chest pain, fever, cough, palpitations.
- ▼ **inflammation of the nerves (Guillain-Barré syndrome):** Pain, weakness, and paralysis in the extremities, numbness or tingling in hands or feet, difficulty with facial movements including speaking, chewing or swallowing.
- ▼ **inflammation of the pancreas (pancreatitis):** abdominal pain, nausea, vomiting, fever
- ▼ **inflammation of the eye:** eye pain, redness and vision changes