

# ULTOMIRIS® PATIENT CARD



## Important Safety Information for Patients Taking ULTOMIRIS® (ravulizumab)

ULTOMIRIS® can lower the ability of your immune system to fight infections, **especially meningococcal infection, which requires immediate medical attention.**



**Get emergency medical care right away if you have any of these signs or symptoms and show this card.**

If you experience any of the following symptoms, you should **immediately call your doctor or seek emergency medical care, preferably in a major emergency medical care center:**

- headache with nausea or vomiting
- headache and fever
- headache with a stiff neck or stiff back
- fever
- fever and rash
- confusion
- muscle aches combined with flu-like symptoms
- eyes sensitive to light

**Keep this card with you at all times** during treatment and for 8 months after your last ULTOMIRIS® dose. Show this card to any healthcare professional involved in your or the child's treatment.

The risk of meningococcal infection may continue for several weeks after the last dose of ULTOMIRIS®.

### Information for Supervising Individuals

Meningitis can become life-threatening within hours.

If you notice, or the child complains about ANY of the following signs or symptoms, seek medical attention immediately. Meningitis and severe blood infection (sepsis) symptoms for infants and children:

#### COMMON SIGNS AND SYMPTOMS:

- Fever, cold hands and feet
- Fretful, dislikes being handled
- Severe muscle pain
- Stiff neck, dislikes bright lights
- Drowsy, floppy, unresponsive
- Severe headache
- Rapid breathing or grunting
- Unusual cry, moaning
- Confusion
- Pale, blotchy skin; spots/rash
- Tense, bulging fontanelle (soft spot)
- Irritability
- Refusing food and vomiting
- Convulsions/seizures

Information for Healthcare Professionals

- This patient has been prescribed ULTOMIRIS® (ravulizumab), which increases the patient’s susceptibility to meningococcal infection (*Neisseria meningitidis*) or other general infections.
- Appropriate vaccination against meningococcal infections should have been administered to the patient at least 2 weeks prior to receiving ULTOMIRIS®. Patients who initiated ULTOMIRIS® less than 2 weeks after receiving meningococcal vaccine must receive prophylactic antibiotics until 2 weeks after vaccination
- Patients must receive vaccination or revaccination according to current national vaccination guidelines for vaccination use.
- Despite the vaccination, the patient may still be susceptible to meningococcal infections or other general infections. Meningococcal infections may become rapidly life-threatening or fatal if not recognized and treated early.
- **Please monitor the emergence of early signs and symptoms of meningococcal infection, evaluate immediately if infection is suspected and treat with appropriate antibiotics if necessary.** The prescribing physician (below) should be contacted as soon as possible.

For more information about ULTOMIRIS®, please refer to the full Singapore Package Insert (PI) or e-mail:  
[MedInfo.SG@astrazeneca.com](mailto:MedInfo.SG@astrazeneca.com).  
In case of safety concerns, call +65 6812 4700 or e-mail:  
[PatientSafety.SG@astrazeneca.com](mailto:PatientSafety.SG@astrazeneca.com).

Patient name \_\_\_\_\_

Parent/Guardian information (if applicable)

\_\_\_\_\_

Hospital where patient is treated \_\_\_\_\_

Physician name \_\_\_\_\_

Tel. number \_\_\_\_\_

Date of Vaccination (DD/MMM/YYYY)	Brand	Serotypes A/C/W/Y